


Advanced Physiology and Pathophysiology Essentials for Clinical Practice 1st Edition Tkacs Test Bank

Chapter 1. THE FOUNDATIONAL CONCEPTS OF CLINICAL PRACTICE

1.	The characteristic, localized cardinal signs of acute inflammation include:
A)	fever.
B)	fatigue.
C)	redness.
D)	granuloma.

2.	The vascular, hemodynamic stage of acute inflammation is initiated by momentary vasoconstriction followed by vasodilation that causes localized:
A)	bleeding.
B)	congestion.
C)	pale skin.
D)	coolness.

3.	The cellular stage of acute inflammation is marked by the movement of leukocytes into the area. Which of the following cells arrives early in great numbers?
A)	Basophils
B)	Lymphocytes
C)	Neutrophils
D)	Platelets



4.	The phagocytosis process involves three distinct steps. What is the initial step in the process?
A)	Engulfment
B)	Intracellular killing
C)	Antigen margination
D)	Recognition and adherence

5.	Which of the following mediators of inflammation causes increased capillary permeability and pain?
A)	Serotonin
B)	Histamine
C)	Bradykinin
D)	Nitric oxide

6.	Inflammatory exudates are a combination of several types. Which of the following exudates is composed of enmeshed necrotic cells?
A)	Serous
B)	Fibrinous
C)	Suppurative
D)	Membranous

7.	The acute-phase systemic response usually begins within hours of the onset of inflammation and includes:
A)	fever and lethargy.
B)	decreased C-reactive protein.
C)	positive nitrogen balance.
D)	low erythrocyte sedimentation rate.

8.	In contrast to acute inflammation, chronic inflammation is characterized by which of the following phenomena?
----	---

A)	Profuse fibrinous exudation
B)	A shift to the left of granulocytes
C)	Metabolic and respiratory alkalosis
D)	Lymphocytosis and activated macrophages

9.	Exogenous pyrogens (interleukin-1) and the presence of bacteria in the blood lead to the release of endogenous pyrogens that:
A)	stabilize thermal control in the brain.
B)	produce leukocytosis and anorexia.
C)	block viral replication in cells.
D)	inhibit prostaglandin release.

10.	An older adult patient has just sheared the skin on her elbow while attempted to boost herself up in bed, an event that has precipitated acute inflammation in the region surrounding the wound. Which of the following events will occur during the vascular stage of the patients inflammation?
A)	Outpouring of exudate into interstitial spaces
B)	Chemotaxis
C)	Accumulation of leukocytes along the epithelium
D)	Phagocytosis of cellular debris

11.	Which of the following individuals most likely has the highest risk of experiencing chronic inflammation?
A)	A patient who has recently been diagnosed with type 2 diabetes
B)	A patient who is a carrier of an antibiotic-resistant organism
C)	A patient who is taking oral antibiotics for an upper respiratory infection
D)	A patient who is morbidly obese and who has a sedentary lifestyle

12.	Which of the following core body temperatures is within normal range?
A)	35.9C (96.6F)

B)	38.0C (100.4F)
C)	35.5C (95.9F)
D)	37.3C (99.1F)

13 .	A postsurgical patient who is recovering in the postanesthetic recovery unit states that she is freezing cold. Which of the following measures is likely to be initiated in the patients hypothalamus in an effort to reduce heat loss?
A)	Opening of arteriovenous (AV) shunts
B)	Reduced exhalation of warmed air
C)	Contraction of pilomotor muscles
D)	Decreased urine production

14 .	An elderly patient is dressed only in a hospital gown and complains of a draft in her room. Consequently, she has requested a warm blanket while she sits in her wheelchair. Which of the following mechanisms of heat loss is most likely the primary cause of her request?
A)	Evaporation and conduction
B)	Radiation and convection
C)	Conduction and convection
D)	Convection and evaporation

15 .	Which of the following pathophysiologic processes are capable of inducing the production of pyrogens? Select all that apply.
A)	Acute inflammation
B)	Obesity
C)	Myocardial infarction
D)	Malignancy
E)	Renal failure

16 .	Which of the following patients is most likely to be susceptible to developing a neurogenic fever?
------	--

A)	A patient who has stage II Alzheimer disease
B)	A patient who has sustained a head injury in a bicycle crash
C)	A patient who has become delirious after the administration of a benzodiazepine
D)	A patient who has begun taking a selective serotonin-reuptake inhibitor (SSRI) for the treatment of depression

17	Patients are commonly administered antipyretics when their oral temperature exceeds 37.5C (99.5F). Which of the following statements related to the rationale for this action is most accurate?
A)	Temperatures in excess of 37.5C (99.5F) can result in seizure activity.
B)	Lower temperatures inhibit the protein synthesis of bacteria.
C)	There is little empirical evidence for this treatment modality.
D)	Most common antipyretics have been shown to have little effect on core temperature.

18	A patient has sought care because of recent malaise and high fever. Upon assessment, the patient states that his current fever began two days earlier, although he states that for the last 2 weeks he is in a cycle of high fever for a couple of days followed by a day or two of normal temperature. Which of the following fever patterns is this patient experiencing?
A)	Recurrent fever
B)	Remittent fever
C)	Sustained fever
D)	Intermittent fever

19	A febrile, 3-week-old infant has been brought to the emergency department by his parents and is currently undergoing a diagnostic workup to determine the cause of his fever. Which of the following statements best conveys the rationale for this careful examination?
A)	The immature hypothalamus is unable to perform normal thermoregulation.

B)	Infants are susceptible to serious infections because of their decreased immune function.
C)	Commonly used antipyretics often have no effect on the core temperature of infants.
D)	Fever in neonates is often evidence of a congenital disorder rather than an infection.

20 .	An 84-year-old patient's blood cultures have come back positive, despite the fact that his oral temperature has remained within normal range. Which of the following phenomena underlies the alterations in fever response that occur in the elderly?
A)	Disturbance in the functioning of the thermoregulatory center
B)	Increased heat loss by evaporation
C)	The presence of comorbidities that are associated with lowered core temperature
D)	Persistent closure of arteriovenous shunts

Answer Key

1.	C
2.	B
3.	C
4.	D
5.	C
6.	D
7.	A
8.	D
9.	B
10 .	A
11 .	D

12	D
.	
13	C
.	
14	B
.	
15	A, C, D
.	
16	B
.	
17	C
.	
18	D
.	
19	B
.	
20	A
.	

Chapter 2. CHEMICAL AND BIOCHEMICAL FOUNDATIONS

1.	Ischemia and other toxic injuries increase the accumulation of intracellular calcium as a result of:
A)	release of stored calcium from the mitochondria.
B)	improved intracellular volume regulation.
C)	decreased influx across the cell membrane.
D)	attraction of calcium to fatty infiltrates.

2.	The patient is found to have liver disease, resulting in the removal of a lobe of his liver. Adaptation to the reduced size of the liver leads to _____ of the remaining liver cells.
----	---

A)	metaplasia
B)	organ atrophy
C)	compensatory hyperplasia
D)	physiologic hypertrophy

3.	A person eating peanuts starts choking and collapses. His airway obstruction is partially cleared, but he remains hypoxic until he reaches the hospital. The prolonged cell hypoxia caused a cerebral infarction and resulting _____ in the brain.
A)	caspase activation
B)	coagulation necrosis
C)	rapid phagocytosis
D)	protein p53 deficiency

4.	Bacteria and viruses cause cell damage by _____, which is unique from the intracellular damage caused by other injurious agents.
A)	disrupting the sodium/potassium ATPase pump
B)	interrupting oxidative metabolism processes
C)	replicating and producing continued injury
D)	decreasing protein synthesis and function

5.	The patient has a prolonged interruption in arterial blood flow to his left kidney, causing hypoxic cell injury and the release of free radicals. Free radicals damage cells by:
A)	destroying phospholipids in the cell membrane.
B)	altering the immune response of the cell.
C)	disrupting calcium storage in the cell.
D)	inactivation of enzymes and mitochondria.

6.	Injured cells have impaired flow of substances through the cell membrane as a result of:
----	--

A)	increased fat load.
B)	altered permeability.
C)	altered glucose utilization.
D)	increased surface receptors.

7.	Reversible adaptive intracellular responses are initiated by:
A)	stimulus overload.
B)	genetic mutations.
C)	chemical messengers.
D)	mitochondrial DNA.

8.	Injured cells become very swollen as a result of:
A)	increased cell protein synthesis.
B)	altered cell volume regulation.
C)	passive entry of potassium into the cell.
D)	bleb formation in the plasma membrane.

9.	A diabetic patient has impaired sensation, circulation, and oxygenation of his feet. He steps on a piece of glass, the wound does not heal, and the area tissue becomes necrotic. The necrotic cell death is characterized by:
A)	rapid apoptosis.
B)	cellular rupture.
C)	shrinkage and collapse.
D)	chronic inflammation.

10.	A 99-year-old woman has experienced the decline of cell function associated with age. A group of theories of cellular aging focus on programmed:
A)	changes with genetic influences.
B)	elimination of cell receptor sites.
C)	insufficient telomerase enzyme.

D)	DNA mutation or faulty repair.
----	--------------------------------

11	An 89-year-old female patient has experienced significant decreases in her mobility and stamina during a 3-week hospital stay for the treatment of a femoral head fracture. Which of the following phenomena most likely accounts for the patients decrease in muscle function that underlies her reduced mobility?
A)	Impaired muscle cell metabolism resulting from metaplasia
B)	Dysplasia as a consequence of inflammation during bone remodeling
C)	Disuse atrophy of muscle cells during a prolonged period of immobility
D)	Ischemic atrophy resulting from vascular changes while on bedrest

12	A 20-year-old college student has presented to her campus medical clinic for a scheduled Papanicolaou (Pap) smear. The clinician who will interpret the smear will examine cell samples for evidence of:
A)	changes in cell shape, size, and organization.
B)	the presence of unexpected cell types.
C)	ischemic changes in cell samples.
D)	abnormally high numbers of cells in a specified field.

13	Which of the following pathophysiologic processes is most likely to result in metastatic calcification?
A)	Benign prostatic hyperplasia
B)	Liver cirrhosis
C)	Impaired glycogen metabolism
D)	Hyperparathyroidism

14	Despite the low levels of radiation used in contemporary radiologic imaging, a radiology technician is aware of the need to minimize her exposure to ionizing radiation. What is the primary rationale for the technicians precautions?
A)	Radiation stimulates pathologic cell hypertrophy and hyperplasia.