


# Lehne's Pharmacology for Nursing Care, 11th Edition Test Bank

by Jacqueline Burchum, Laura Rosenthal



## Table Of Contents

### **UNIT I: INTRODUCTION**

- Chapter 1. Orientation to Pharmacology
- Chapter 2. Application of Pharmacology in Nursing Practice
- Chapter 3. Drug Regulation, Development, Names, and Information

### **UNIT II: BASIC PRINCIPLES OF PHARMACOLOGY**

- Chapter 4. Pharmacokinetics
- Chapter 5. Pharmacodynamics
- Chapter 6. Drug Interactions
- Chapter 7. Adverse Drug Reactions and Medication Errors
- Chapter 8. Individual Variation to Drug Responses
- Chapter 9. Genetic and Genomic Considerations NEW!
- Chapter 10. Introduction to Immunomodulators NEW!

### **UNIT III: DRUG THERAPY ACROSS THE LIFE SPAN**

- Chapter 11. Drug Therapy During Pregnancy and Breast-Feeding
- Chapter 12. Drug Therapy in Pediatric Patients
- Chapter 13. Drug Therapy in Geriatric Patients

### **UNIT IV: CENTRAL NERVOUS SYSTEM DRUGS**

#### *Section 1: Introduction*

- Chapter 14. Basic Principles of Neuropharmacology
- Chapter 15. Physiology of the Peripheral Nervous System

#### *Section 2: Cholinergic Drugs*

- Chapter 16. Muscarinic Agonists
- Chapter 17. Muscarinic Antagonists NEW!
- Chapter 18. Cholinesterase Inhibitors and Their Use in Myasthenia Gravis
- Chapter 19. Drugs That Block Nicotinic Cholinergic Transmission: Neuromuscular Blocking Agents

#### *Section 3: Adrenergic Drugs*

- Chapter 20. Adrenergic Agonists
- Chapter 21. Adrenergic Antagonists
- Chapter 22. Indirect-Acting Antiadrenergic Agents

### **UNIT V: CENTRAL NERVOUS SYSTEM DRUGS**

#### *Section 4: Introduction*

- Chapter 23. Introduction to Central Nervous System Pharmacology
- Section 5: Drugs for Neurodegenerative Disorders
- Chapter 24. Drugs for Parkinson's Disease

Chapter 25. Drugs for Alzheimer's Disease

Chapter 26. Drugs for Multiple Sclerosis

*Section 6: Neurologic Drugs*

Chapter 27. Drugs for Epilepsy

Chapter 28. Drugs for Muscle Spasm and Spasticity

*Section 7: Drugs for Pain*

Chapter 29. Local Anesthetics

Chapter 30. General Anesthetics

Chapter 31. Opioid Analgesics, Opioid Antagonists, and Nonopioid Centrally Acting Analgesics

Chapter 32. Pain Management in Patients with Cancer

Chapter 33. Drugs for Headache

*Section 8: Psychotherapeutic Drugs*

Chapter 34. Antipsychotic Agents and Their Use in Schizophrenia

Chapter 35. Antidepressants

Chapter 36. Drugs for Bipolar Disorder

Chapter 37. Sedative-Hypnotic Drugs

Chapter 38. Management of Anxiety Disorders

Chapter 39. Central Nervous System Stimulants and Attention-Deficit/Hyperactivity Disorder

*Section 9: Substance Use Disorders*

Chapter 40. Drug Abuse I: Basic Considerations

Chapter 41. Drug Abuse II: Alcohol

Chapter 42. Drug Abuse III: Nicotine and Smoking

Chapter 43. Drug Abuse IV: Major Drugs of Abuse Other Than Alcohol and Nicotine

**UNIT VI: DRUGS THAT AFFECT FLUID AND ELECTROLYTE BALANCE**

Chapter 44. Diuretics

Chapter 45. Agents Affecting the Volume and Ion Content of Body Fluids

**UNIT VII: DRUGS THAT AFFECT THE HEART, BLOOD VESSELS, AND BLOOD**

Chapter 46. Review of Hemodynamics

Chapter 47. Drugs Acting on the Renin-Angiotensin-Aldosterone System

Chapter 48. Calcium Channel Blockers

Chapter 49. Vasodilators

Chapter 50. Drugs for Hypertension

Chapter 51. Drugs for Heart Failure

Chapter 52. Antidysrhythmic Drugs

Chapter 53. Prophylaxis of Atherosclerotic Cardiovascular Disease: Drugs That Help Normalize Cholesterol and Triglyceride Levels

Chapter 54. Drugs for Angina Pectoris

Chapter 55. Anticoagulant, Antiplatelet, and Thrombolytic Drugs

Chapter 56. Management of ST-Elevation Myocardial Infarction

Chapter 57. Drugs for Hemophilia

Chapter 58. Drugs for Deficiency Anemias

Chapter 59. Hematopoietic Agents

**UNIT VIII: DRUGS FOR ENDOCRINE DISORDERS**

Chapter 60. Drugs for Diabetes Mellitus

Chapter 61. Drugs for Thyroid Disorders

Chapter 62. Drugs Related to Hypothalamic and Pituitary Function

Chapter 63. Drugs for Disorders of the Adrenal Cortex

**UNIT IX: WOMEN'S HEALTH**

Chapter 64. Estrogens and Progestins: Basic Pharmacology and Noncontraceptive Applications

Chapter 65. Birth Control

Chapter 66. Drug Therapy of Infertility

Chapter 67. Drugs That Affect Uterine Function

**UNIT X: MEN'S HEALTH**

Chapter 68. Androgens

Chapter 69. Drugs for Erectile Dysfunction and Benign Prostatic Hyperplasia

**UNIT XI: ANTI-INFLAMMATORY, ANTIALLERGIC, AND IMMUNOLOGIC DRUGS**

Chapter 70. Review of the Immune System

Chapter 71. Childhood Immunization

Chapter 72. Immunosuppressants

Chapter 73. Antihistamines

Chapter 74. Cyclooxygenase Inhibitors: Nonsteroidal Anti-inflammatory Drugs and Acetaminophen

Chapter 75. Glucocorticoids in Nonendocrine Disorders

**UNIT XII: DRUGS FOR BONE AND JOINT DISORDERS**

Chapter 76. Drug Therapy of Rheumatoid Arthritis

Chapter 77. Drug Therapy of Gout

Chapter 78. Drugs Affecting Calcium Levels and Bone Mineralization

**UNIT XIII: RESPIRATORY TRACT DRUGS**

Chapter 79. Drugs for Asthma and Chronic Obstructive Pulmonary Disease

Chapter 80. Drugs for Allergic Rhinitis, Cough, and Colds

**UNIT XIV: GASTROINTESTINAL DRUGS**

Chapter 81. Drugs for Peptic Ulcer Disease

Chapter 82. Laxatives

Chapter 83. Other Gastrointestinal Drugs

**UNIT XV: NUTRITION AND COMPLIMENTARY THERAPY**

Chapter 84. Vitamins

Chapter 85. Drugs for Weight Loss

Chapter 86. Complementary and Alternative Therapy

**UNIT XVI: THERAPY OF INFECTIOUS DISEASES**

Chapter 87. Basic Principles of Antimicrobial Therapy

Chapter 88. Drugs That Weaken the Bacterial Cell Wall I: Penicillins

Chapter 89. Drugs That Weaken the Bacterial Cell Wall II: Cephalosporins, Carbapenems, Vancomycin, Telavancin, Aztreonam, Teicoplanin, and Fosfomycin

Chapter 90. Bacteriostatic Inhibitors of Protein Synthesis: Tetracyclines, Macrolides, and Others

Chapter 91. Aminoglycosides: Bactericidal Inhibitors of Protein Synthesis

Chapter 92. Sulfonamides and Trimethoprim

Chapter 93. Drug Therapy of Urinary Tract Infections

Chapter 94. Antimycobacterial Agents: Drugs for Tuberculosis, Leprosy, and Mycobacterium avium Complex Infection

Chapter 95. Miscellaneous Antibacterial Drugs: Fluoroquinolones, Metronidazole, Daptomycin, Rifampin, Rifaximin, Bacitracin, and Polymyxins

Chapter 96. Antifungal Agents

Chapter 97. Antiviral Agents I: Drugs for Non-HIV Viral Infections

Chapter 98. Antiviral Agents II: Drugs for HIV Infection and Related Opportunistic Infections

Chapter 99. Drug Therapy of Sexually Transmitted Diseases

Chapter 100. Antiseptics and Disinfectants

**UNIT XVII: CHEMOTHERAPY OF PARASITIC DISEASES**

Chapter 101. Anthelmintics

Chapter 102. Antiprotozoal Drugs I: Antimalarial Agents

Chapter 103. Antiprotozoal Drugs II: Miscellaneous Agents

Chapter 104. Ectoparasiticides

**UNIT XVIII: CANCER CHEMOTHERAPY**

Chapter 105. Basic Principles of Cancer Chemotherapy

Chapter 106. Anticancer Drugs I: Cytotoxic Agents

Chapter 107. Anticancer Drugs II: Noncytotoxic Agents

**UNIT XIX: MISCELLANEOUS DRUGS AND THERAPIES**

Chapter 108. Drugs for the Eye

Chapter 109. Drugs for the Skin

Chapter 110. Drugs for the Ear

**UNIT XX: TOXICOLOGY**

Chapter 111. Management of Poisoning

Chapter 112. Potential Weapons of Biologic, Radiologic, and Chemical Terrorism

## Chapter 01: Orientation to Pharmacology

### Burchum: Lehne's Pharmacology for Nursing Care, 11th Edition

---

#### MULTIPLE CHOICE

1. The nurse is teaching a patient how a medication works to treat an illness. To do this, the nurse will rely on knowledge of which topic?
  - a. Clinical pharmacology
  - b. Drug efficacy
  - c. Pharmacokinetics
  - d. Pharmacotherapeutics

ANS: D

Pharmacotherapeutics is the study of the use of drugs to diagnose, treat, and prevent conditions. Clinical pharmacology is concerned with all aspects of drug-human interactions. Drug efficacy measures the extent to which a given drug causes an intended effect. Pharmacokinetics is the study of the impact of the body on a drug.

PTS: 1

2. What is a desired outcome when a drug is described as easy to administer?
  - a. It can be stored indefinitely without need for refrigeration.
  - b. It does not interact significantly with other medications.
  - c. It enhances patient adherence to the drug regimen.
  - d. It is usually relatively inexpensive to produce.

ANS: C

A major benefit of drugs that are easy to administer is that patients taking them are more likely to comply with the drug regimen. Drugs that are easy to give may have the other attributes listed, but those properties are independent of ease of administration.

PTS: 1

3. A patient tells the nurse that an analgesic he will begin taking may cause drowsiness and will decrease pain up to 4 hours at a time. Based on this understanding of the drug's effects by the patient, the nurse will anticipate which outcome?
  - a. Decreased chance of having a placebo effect
  - b. Decreased motivation to take the drug
  - c. Improved compliance with the drug regimen
  - d. Increased likelihood of drug overdose

ANS: C

A drug is effective if it produces the intended effects, even if it also produces side effects. Patients who understand both the risks and benefits of taking a medication are more likely to comply with the drug regimen.

PTS: 1

## MULTIPLE RESPONSE

1. What are considered the 'Big Three' properties of an ideal drug? (*Select all that apply.*)
  - a. Irreversible action
  - b. Effectiveness
  - c. Safety
  - d. Selectivity
  - e. A recognizable trade name

ANS: B, C, D

The 'Big Three' properties of the ideal drug are effectiveness, safety, and selectivity.

PTS: 1

2. Before administering a medication, what does the nurse need to know to evaluate how individual patient variability might affect the patient's response to the medication? (*Select all that apply.*)
  - a. Chemical stability of the medication
  - b. Ease of administration
  - c. Family medical history
  - d. Patient's age
  - e. Patient's diagnosis

ANS: C, D, E

The family medical history can indicate genetic factors that may affect a patient's response to a medication. Patients of different ages can respond differently to medications. The patient's illness can affect how drugs are metabolized. The chemical stability of the medication and the ease of administration are properties of drugs.

PTS: 1

## Chapter 02: Application of Pharmacology in Nursing Practice

### Burchum: Lehne's Pharmacology for Nursing Care, 11th Edition

---

## MULTIPLE CHOICE

1. A patient is using a metered-dose inhaler containing albuterol for asthma. The medication label instructs the patient to administer "2 puffs every 4 hours as needed for coughing or wheezing." The patient reports feeling jittery sometimes when taking the medication, and doesn't feel that the medication is always effective. Which action is outside the nurse's scope of practice?
  - a. Asking the patient to demonstrate the use of the inhaler
  - b. Assessing the patient's exposure to tobacco smoke
  - c. Auscultating lung sounds and obtaining vital signs
  - d. Suggesting that the patient use 1 puff to reduce side effects

ANS: D

It is not within the nurse's scope of practice to change the dose of a medication without an order from a prescriber. Asking the patient to demonstrate inhaler use helps the nurse to evaluate the patient's ability to administer the medication properly and is part of the nurse's evaluation. Assessing tobacco smoke exposure helps the nurse determine whether nondrug therapies, such as smoke avoidance, can be used as an adjunct to drug therapy. Performing a physical assessment helps the nurse evaluate the patient's response to the medication.

PTS: 1

2. A postoperative patient is being discharged home with acetaminophen/hydrocodone [Norco] for pain. The patient asks the nurse about using Tylenol for fever. Which statement by the nurse is correct?
  - a. "It is not safe to take over-the-counter drugs with prescription medications."
  - b. "Taking the two medications together poses a risk of drug toxicity."
  - c. "There are no known drug interactions, so this will be safe."
  - d. "Tylenol and Norco are different drugs, so there is no risk of overdose."

ANS: B

Tylenol is the trade name and acetaminophen is the generic name for the same medication. It is important to teach patients to be aware of the different names for the same drug to minimize the risk of overdose. Over-the-counter (OTC) medications and prescription medications may be taken together unless significant harmful drug interactions are possible. Even though no drug interactions are at play in this case, both drugs contain acetaminophen, which could lead to toxicity.

PTS: 1

3. The nurse is preparing to care for a patient who will be taking an antihypertensive medication. Which action by the nurse is part of the assessment step of the nursing process?
  - a. Asking the prescriber for an order to monitor serum drug levels
  - b. Monitoring the patient for drug interactions after giving the medication
  - c. Questioning the patient about over-the-counter medications
  - d. Taking the patient's blood pressure throughout the course of treatment

ANS: C

The assessment part of the nursing process involves gathering information before beginning treatment, and this includes asking about other medications the patient may be taking. Monitoring serum drug levels, watching for drug interactions, and checking vital signs after giving the medication are all part of the evaluation phase.

PTS: 1

4. A postoperative patient reports pain, which the patient rates as an 8 on a scale from 1 to 10 (10 being the most extreme pain). The prescriber has ordered acetaminophen [Tylenol] 650 mg PO every 6 hours PRN pain. What will the nurse do?
  - a. Ask the patient what medications have helped with pain in the past.
  - b. Contact the provider to request a different analgesic medication.
  - c. Give the pain medication and reposition the patient to promote comfort.

- d. Request an order to administer the medication every 4 hours.

ANS: B

The nursing diagnosis for this patient is severe pain. Acetaminophen is given for mild to moderate pain, so the nurse should ask the prescriber to order a stronger analgesic medication. Asking the patient to tell the nurse what has helped in the past is part of an initial assessment and should be done preoperatively and not when the patient is having severe pain. Because the patient is having severe pain, acetaminophen combined with nondrug therapies will not be sufficient. Increasing the frequency of the dose of a medication for mild pain will not be effective.

PTS: 1

5. A patient newly diagnosed with diabetes is to be discharged from the hospital. The nurse teaching this patient about home management should begin by doing what?
- Asking the patient to demonstrate how to measure and administer insulin
  - Discussing methods of storing insulin and discarding syringes
  - Giving information about how diet and exercise affect insulin requirements
  - Teaching the patient about the long-term consequences of poor diabetes control

ANS: A

Because insulin must be given correctly to control symptoms and because an overdose can be fatal, it is most important for the patient to know how to administer it. Asking for a demonstration of technique is the best way to determine whether the patient has understood the teaching. When a patient is receiving a lot of new information, the information presented first is the most likely to be remembered. The other teaching points are important as well, but they are not as critical and can be taught later.

PTS: 1

6. The nurse receives an order to give morphine 5 mg IV every 2 hours PRN pain. Which action is *not* part of the six rights of drug administration?
- Assessing the patient's pain level 15 to 30 minutes after giving the medication
  - Checking the medication administration record to see when the last dose was administered
  - Consulting a drug manual to determine whether the amount the prescriber ordered is appropriate
  - Documenting the reason the medication was given in the patient's electronic medical record

ANS: A

Assessing the patient's pain after administering the medication is an important part of the nursing process when giving medications, but it is not part of the six rights of drug administration. Checking to see when the last dose was given helps ensure that the medication is given at the right time. Consulting a drug manual helps ensure that the medication is given in the right dose. Documenting the reason for a pain medication is an important part of the right documentation—the sixth right.

PTS: 1

7. A patient tells a nurse that a medication prescribed for recurrent migraine headaches is not working. What will the nurse do?
- Ask the patient about the number and frequency of tablets taken.
  - Assess the patient's headache pain on a scale from 1 to 10.
  - Report the patient's complaint to the prescriber.
  - Suggest biofeedback as an adjunct to drug therapy.

ANS: A

When evaluating the effectiveness of a drug, it is important to determine whether the patient is using the drug as ordered. Asking the patient to tell the nurse how many tablets are taken and how often helps the nurse determine compliance. Assessing current pain does not yield information about how well the medication is working unless the patient is currently taking it. The nurse should gather as much information about compliance, symptoms, and drug effectiveness as possible before contacting the prescriber. Biofeedback may be an effective adjunct to treatment, but it should not be recommended without complete information about drug effectiveness.

PTS: 1

8. A nurse is preparing to administer medications. Which patient would the nurse consider to have the greatest predisposition to an adverse reaction?
- A 30-year-old man with kidney disease
  - A 75-year-old woman with cystitis
  - A 50-year-old man with an upper respiratory tract infection
  - A 9-year-old boy with an ear infection

ANS: A

The individual with impaired kidney function would be at risk of having the drug accumulate to a toxic level because of potential excretion difficulties. Cystitis is an infection of the bladder and not usually the cause of excretion problems that might lead to an adverse reaction from a medication. A respiratory tract infection would not predispose a patient to an adverse reaction, because drugs are not metabolized or excreted by the lungs. A 9-year-old boy would not have the greatest predisposition to an adverse reaction simply because he is a child; nor does an ear infection put him at greater risk.

PTS: 1

9. A nurse consults a drug manual before giving a medication to an 80-year-old patient. The manual states that older adult patients are at increased risk for hepatic side effects. Which action by the nurse is correct?
- Contact the provider to discuss an order for pretreatment laboratory work.
  - Ensure that the drug is given in the correct dose at the correct time to minimize the risk of adverse effects.
  - Notify the provider that this drug is contraindicated for this patient.
  - Request an order to give the medication intravenously so that the drug does not pass through the liver.

ANS: A



The drug manual indicates that this drug should be given with caution to elderly patients. Getting information about liver function before giving the drug establishes baseline data that can be compared with post-treatment data to determine whether the drug is affecting the liver. Giving the correct dose at the correct interval helps to minimize risk, but without baseline information, the effects cannot be determined. The drug is not contraindicated.

PTS: 1

10. A patient has been receiving intravenous penicillin for pneumonia for several days and begins to complain of generalized itching. The nurse auscultates bilateral wheezing and notes a temperature of 38.5°C (101°F). Which is the correct action by the nurse?
- Administer the next dose and continue to evaluate the patient's symptoms.
  - Ask the prescriber if an antihistamine can be given to relieve the itching.
  - Contact the prescriber to request an order for a chest radiograph.
  - Hold the next dose and notify the prescriber of the symptoms.

ANS: D

Pruritus and wheezing are signs of a possible allergic reaction, which can be fatal; therefore, the medication should not be given and the prescriber should be notified. When patients are having a potentially serious reaction to a medication, the nurse should not continue giving the medication. Antihistamines may help the symptoms of an allergic reaction, but the first priority is to stop the medication. Obtaining a chest radiograph is not helpful.

PTS: 1

11. A postoperative patient has orders for morphine sulfate 1 to 2 mg IV every 1 hour PRN for severe pain and acetaminophen-hydrocodone [Norco] 5 mg PO every 4 to 6 hours PRN for moderate pain. The patient reports pain at a level of 8 on a scale of 1 to 10, with 10 being the worst pain. Which action by the nurse is appropriate?
- Administer acetaminophen-hydrocodone 5 mg PO every 4 hours.
  - Administer acetaminophen-hydrocodone 5 mg PO every 6 hours and change to every 4 hours if not effective.
  - Administer morphine sulfate 1 mg IV every 1 hour until pain subsides.
  - Administer morphine sulfate 2 mg IV and evaluate the patient's pain in 15 to 30 minutes.

ANS: D

With PRN medications, the schedule is not fixed and the administration of these medications depends on the patient's condition. It is the nurse's responsibility to assess the patient's condition and then give the appropriate PRN medication. In this case, the patient has severe pain and should receive MS IV. Either 1 or 2 mg may be given, but the nurse must evaluate the effectiveness of the pain medication within 15 to 30 minutes to help determine subsequent doses. Acetaminophen-hydrocodone is not appropriate because it is ordered for moderate pain and this patient reports severe pain. Giving MS IV every hour is not appropriate for a PRN medication unless the patient's condition warrants it.

PTS: 1

12. The nurse is teaching a patient about home administration of insulin to treat diabetes mellitus. As part of the teaching, the patient and nurse identify goals to maintain specific blood glucose ranges. This represents which aspect of the nursing process?
- Assessment
  - Evaluation
  - Implementation
  - Planning

ANS: D

In the planning step, the nurse delineates specific interventions directed at solving or preventing problems. When creating the care plan, the nurse defines goals, sets priorities, and establishes criteria for evaluating success. The assessment step involves collecting data about the patient. The evaluation step involves evaluating the medication effectiveness. The implementation step identifies actions that are taken to administer the drug.

PTS: 1

#### **MULTIPLE RESPONSE**

1. A nurse is reviewing a patient's medical record before administering a medication. Which factors can alter the patient's physiologic response to the drug? (*Select all that apply.*)
- Ability to swallow pills
  - Age
  - Genetic factors
  - Gender
  - Height

ANS: B, C, D

Age, genetic factors, and gender influence an individual patient's ability to absorb, metabolize, and excrete drugs; therefore, these factors must be assessed before a medication is administered. A patient's ability to swallow pills, although it may determine the way a drug is administered, does not affect the physiologic response. Height does not affect response; weight and the distribution of adipose tissue can affect the distribution of drugs.

PTS: 1

#### **Chapter 03: Drug Regulation, Development, Names, and Information** **Burchum: Lehne's Pharmacology for Nursing Care, 11th Edition**

---

#### **MULTIPLE CHOICE**

1. A nurse educator is conducting a continuing education class on pharmacology. To evaluate the learning of the nurses in the class, the nurse educator asks, "Which drug name gives information about the drug's pharmacologic classification?" Which is the correct response?
- Amoxicillin
  - Tylenol
  - Cipro
  - Motrin